

## “Comparative evaluation of Sexually Transmitted and Non Sexually Transmitted, Reproductive tract infection at tertiary care centre in western Rajasthan, India”

Priyanka Soni<sup>1,\*</sup>, Smita Kulsrestha<sup>2</sup>, R.S. Parihar<sup>3</sup>, P.K. Khatri<sup>4</sup>

<sup>1</sup>PG Student, <sup>2</sup>Associate Professor, <sup>3</sup>Professor, <sup>4</sup>Professor & Head, Dept. of Microbiology, Dr. S.N. Medical College, Jodhpur

**\*Corresponding Author:**

Email: priyanka.soni88@gmail.com

### Abstract

**Introduction:** Reproductive Tract infections among women of reproductive age group (including both sexually Transmitted Infection (STI) & Non STI) are responsible for major ill health and have important concern as these are associated with risk of HIV Transmission & complications like Pelvic inflammatory Disease (PID), Infertility etc. This study was conducted to determine prevalence of microorganism & their antibiogram & their association with socio demographic profile & menstrual hygiene.

**Methods:** This was a cross sectional study included 200 women with discharge of 15-44 year age. Detail history was taken about their socio demographic data, Symptoms & menstrual hygiene; vaginal or endo cervical swab & blood sample (for serological diagnosis of syphilis) were collected.

**Results:** Among STI most common infection is Candidiasis (45.6%) & Bacterial Vaginosis (27.8%) whereas among Non STI Esch. coli (27.3%) & Staph. aureus (26.4%). Among STI most common symptom was lower abdominal Pain & Lower backache whereas in Non-STI Vulvular itching & fever. Both STI & Non STI shows statically significant positive association with Labourer population & who uses cloth during their menstrual period whereas STI also shows statically significant positive association with younger, lower education & lower socio economic status population. Staph. aureus highly sensitive to linezolid, vancomycin whereas Esch.coli was most common gram negative bacteria & highly sensitive to imipenem & piperacillin-tazobactam.

**Conclusions:** To conclude, by proper assessment of Prevalence & risk factor of STI and Non STI & by health education, we can reduce these infection & antibiotic resistance by prudent use of antibiotics.

**Key Words:** Reproductive tract infection, STI, Non-STI

Access this article online	
Quick Response Code:	Website: www.innovativepublication.com
	DOI: 10.5958/2394-5478.2016.00025.X

### Introduction

Reproductive tract infections (RTI) are a global health problem among women, living in South East Asian Region (SEAR) countries. Studies have found the prevalence of RTI in India, Bangladesh, Egypt, and Kenya is in the range of 52–90 per cent. More than a million women and infants die of the complications of RTI every year.<sup>1</sup>

The vaginal flora is a complicated environment, containing dozens of microbiological species in variable quantities and relative proportions. A complex and intricate balance of microorganisms maintains the normal vaginal flora. It is mainly dominated by members of the genus *Lactobacillus*, which maintains the generally acidic vaginal pH.<sup>2</sup> The frequent cause of vaginal discharge is an infection or colonization with different microorganisms.<sup>3</sup> Vaginitis, whether infectious or not, poses one of the most common problems in gynecology, and is one of the main reasons leading the females to seek advice from a physician approximately

10 million office visits annually.<sup>4-7</sup> However, diagnosis and treatment can be elusive, if based on clinical symptoms and the characterization of vaginal discharge alone, leading to a lack of relief from the symptoms.<sup>8-9</sup>

A variety of factors that put women at risk of reproductive tract infection as well as consequences for women arising from such morbidity, such as socio-economic, demographic, sexual, medical, behavioral practices, personal hygiene behavior have not been adequately explored in India. Concerted efforts are needed to provide useful information to health planners and policymakers so that appropriate strategies can be designed to bring about an improvement in reproductive health of women.<sup>10</sup>

This study was conducted to determine Comparative evaluation of STI & Non-STI using following parameters- (1) Prevalence of microorganism causing STI & Non-STI.<sup>(2)</sup> Antibiotic sensitivity pattern of aerobic vaginal pathogen (2) Comparative study of frequency of Symptoms (3) Association of RTI with education status, socioeconomic status, Occupation, Rural/Urban & Menstrual hygiene among women of reproductive age group.

### Materials and Methods

**Study Design:** It is a hospital based cross-sectional Study conducted in Department of Microbiology. Dr. S.

N. Medical College and associated group of hospital-Umaid Hospital, Jodhpur from January 2014 to August 2014.

**Study Population:** 200 women in age group 15-44 year were selected, who had abnormal vaginal discharge both from indoor and outdoor.

**Study Tools:** A predesigned, pretested, and semi structured questionnaire was used to take the interview of eligible women. The questionnaire had both open and closed ended questions about RTI symptoms, their socio demographic data, occupation and menstrual hygiene.

**Ethical Issues:** The aims, objectives, and procedure of the study were explained to all the women. Informed consent was taken from all the participants. Complete confidentiality regarding patient information was maintained through all the stages of the study.

**Statistical Tests:** Data was analysed using SPSS version 17. Percentage and proportions were calculated for prevalence of the symptoms and socio demographic data. Chi square and Fisher Exact test were used as tests of significance in univariate analysis. A *P* value of less than 0.05 was considered significant.

**Per speculum examination** was carried out and Samples were collected and sent for investigation at Microbiology Laboratory at Umaid Hospital.

- Endocervical swab: gonococcal infection (Gram staining)
- Vaginal swab: 2 vaginal swab from each women, one for wet preparation, KOH mount & pH testing ; and second for Culture & sensitivity and Gram staining
- Blood investigations: RPR (syphilis) and TPHA.

#### Criteria used for laboratory diagnosis

#### Reproductive tract infections: Diagnostic criteria for sexually transmitted infections

- **Gonorrhoea-** Identification of Gm -ve intracellular diplococci in Gram stained cervical smear
- **Trichomonas vaginitis-**  $\geq 1$  actively moving trichomonad (saline wet preparation), pH > 5
- **Bacterial vaginosis-** Presence of "Clue" cells (saline wet preparation and in gram stained smear), Vaginal Ph > 5
- **Candidiasis-** Presence of budding yeast cells and pseudohyphae (KOH wet preparation) and vaginal Ph < 5
- **Syphilis-** Nontreponemal test – RPR, Confirmation by TPHA test

**Limitation of study:** Diagnosing Trichomonas infection by wet mount preparation is approximately 60-98.2% sensitive compared to trichomonas culture.<sup>11-12</sup>

The sensitivity of Clue cells as diagnostic criteria for bacterial vaginosis is 80-98% when compared with more elaborate criteria for determining bacterial vaginosis.<sup>13</sup> Detection of intracellular gram-negative diplococci in endocervical mucus is quite specific but  $\leq$  50% sensitive for gonorrhoea.<sup>14</sup> Diagnosis of

Chlamydia infection, as a causative agent of STI is not studied due to non-availability of resources.

**Reproductive tract infections: Diagnosis of non-sexually transmitted infections:** All vaginal discharge sample were collected with the sterile swabs tested for organism causing STI and inoculated on blood agar, Mac Conkey's agar and Chocolate agar for diagnosis of Non-STI. The aerobically incubated organisms were identified with the help of colonial morphology, gram staining and biochemical analysis.<sup>15</sup> Isolated organisms were subjected to sensitivity testing by Kirby-Bauer disc diffusion method, using CLSI Standards criteria, to interpret diameter of inhibition zone.<sup>16</sup>

#### Results

In this study STI are more common in younger age group {15-24years (50%)( $p < 0.05$ )} whereas Non STI found almost equally in all age groups( $p > 0.54$ ). STI are more frequent in low education status, labourers, rural Population, low socio economic status and in women who uses cloth during their menstrual period ( $p < 0.05$ ). Non-STI infections are significantly more common in Labourers and in women who uses cloth during their menstrual period. ( $p < 0.05$ ) (Table 3)

Majority of women with STI and Non-STI are suffer from abnormal excessive Vaginal discharge (100%); Among STI most common symptom was lower abdominal Pain (71.6%) followed by Lower backache (66.7%) whereas in Non-STI Vulvular itching & burning (65%), fever (60%) & lower abdominal pain (60%) were most common. Lower backache is significantly more in patients of STI ( $p < 0.03$ ) & fever in patient of Non-STI ( $p < 0.0001$ ) (Table 2)

This study shows that among STI most common infection is Candidiasis (45.6%) followed by Bacterial Vaginosis (27.8%), Trichomonas Vaginalis (22.2%), Syphilis (2.2%) and gonorrhoea (2.2%). Among Non-STI most common infection caused by Esch. coli (27.3%), & Staph. aureus (26.4%), Coagulase neg. Staph. (16.1%) followed by Klebsiella sp. (7.2%), Enterobacter sp. (5.6%), Citrobacter sp. (3.6%), Pseudomonas sp. (3%). (Table 1)

The sensitivity pattern of Staph.aureus -100% sensitive to linezolid, >80% to Vancomycin & imipenam, 72% to amoxyclave, and almost 50% to Ampicillin Cefixime, Ceftriaxone & Cefazolin; 72.5% isolates were resistant to methicillin. Fig. 1

Esch.coli was most common gram negative bacteria. GNB found mostly sensitive >80% to imipenam, Piperacillin- Tazobactam, 63-73% sensitive to Amikacin, Gentamicin, Ciprofloxacin & Ofloxacin, 50% to Cefixime & ceftriaxone. (Table 4 and Fig. 2)

#### Discussion

The highest incidence of STI in this study was noted among young, sexually active females, at the two age groups, i.e., 15-24 years (50%) followed by 25-34

years (35.5%); in non-STI % are 31.8% and 38.2% respectively, these similar to study by Khan et al<sup>2</sup> and Shamim et al.<sup>17</sup>

Prevalence of RTI/STI were more in low education and rural areas patients. This finding is comparable to a study carried out in Rajasthan by Bansal et al.<sup>18</sup> The different pattern of infection (STI & Non-STI) in the present study may be owing to the prevalent conditions like health education, sanitation and medical coverage available in each country.

*Escherichia coli* and *Staphylococcus aureus* were commonly isolated vaginal pathogens in my study. According to Shamim et al<sup>17</sup>, *Staphylococcus aureus* was most prevalent followed by *Esch. coli*. Vigneswaran R et al<sup>22</sup> found *E. coli* and group B. *Streptococci* to be the important pathogens associated with mid-trimester pregnancy losses, alongside the classic bacterial vaginosis organisms

The most effective chemotherapeutic agents observed against *Staphylococcus aureus* in this study were linezolid (100%), imipenem (83%), vancomycin (89.6%), lesser activity has been noted against penicillins & cephalosporins (36.8%). In most cases of *Staphylococcus aureus*, resistance to penicillin is attributable to  $\beta$ -lactamase production. Therefore, penicillin in combination with one of the  $\beta$  lactamase inhibitors gives much better results<sup>23-24</sup> clearly seen from this study (Ampicillin 51% & coamoxiclav 72%).


The most effective chemotherapeutic agents against gram-negative rods (*Escherichia coli* and *Klebsiella sp.*) were imipenem (86.7%, 87.5%) and piperacillin/tazobactam (83.4%, 87.5%), whereas the antimicrobials with least affectivity against most of

gram negative bacilli were those belonging to the groups of cephalosporins (50% sensitivity); fluoroquinolones and aminoglycosides have moderate activity(62-75%). This in accordance with the study by Tariq et al<sup>17</sup>, Shamim et al.<sup>20</sup>


Aerobic vaginitis typically does not respond to antibacterial vaginosis medication so has to be treated by antibiotics according to the culture and sensitivity. An optimal treatment scheme for aerobic vaginitis, which includes antibacterial agents and simultaneously normalizing the vaginal ecosystem, has not been established until today. So, there is need for such type of study which diagnose aerobic vaginal pathogen, their sensitivity profile and causative agent of STI for proper treatment of Reproductive tract infection.

### Conclusions

To conclude, the high prevalence of gynaecological infections demands that these patients should be investigated thoroughly for STI and Non-STI simultaneously. As the culture provides the identification of causative microorganisms, it must invariably be done. Currently the antibiotics showing good sensitivity are carbapenems and  $\beta$ -lactams  $\beta$ -lactamase inhibitor combinations, but are very expensive and out of reach of the poor patients, so there is need for an antibiotic policy for its rational use. Proper assessment of prevalence and risk factor of STI and Non STI and their modification by health education & Government policies may help to reduce these infection in near future.



**Fig. 1: Staphylococcus aureus - Antibiotic Sensitivity Pattern**  
**Staph. aureus Antibiotic Sensitivity Pattern**



**Fig. 2: Esch. coli - Antibiotic Sensitivity Pattern**  
Esch. Coli Antibiotic Sensitivity Pattern

**Table 1: Distribution of Patients According to causative agents of STI and Non-STI**

STI		Non STI	
Causative agent	No (%)	Causative agent	No (%)
Trichomonas	20(22.2%)	Esch.coli	30(27.3%)
Syphilis	2(2.2%)	Staph. Aureus	29(26.4%)
Candida sp.	41(45.6%)	Coagulase neg. Staph	18(16.1%)
Bacterial Vaginosis	25(27.8%)	Klebsiella sp.	8(7.2%)
Gonorrhoea	2(2.2%)	Enterobacter	6(5.6%)
		Citrobacter	4(3.6%)
		Pseudomonas	3(2.8%)
		Beta haemolytic Streptococci	2(1.8%)
		Organism not grown	10(9.09%)
Total	90		110

**Table 2: Comparative evaluation of patients of STI and Non-STI According to symptoms**

Symptoms	STI 90		Non STI 110		P value
	No	%	No	%	
Abnormal excessive discharge	90	100%	110	100%	0.920
Lower backache	60	66.7%	57	52%	0.03
Lower Abdominal Pain	64	71.1%	66	60%	0.102
Associated Fever	20	22.2%	66	60%	<0.0001
genital ulceration	2	2.2%	8	7.27%	0.123
Vulvar itching & Burning	52	57.8%	71	65%	0.328
Dysmenorrhea	18	20%	31	28.18%	0.182
Menstrual disturbance	37	41.1%	36	32.72%	0.221

**Table 3 Comparative evaluation of socio demographic factor among patient of STI and Non-STI**

Socio-demographic Factor	STI (90)		Non-STI (110)	
	No.	P value	No.	P value
<b>(A) Age group</b>				
15-24 year	45	<b>0.0002</b>	35	0.543
25-34 year	32		42	
35-44 year	13		33	
<b>(B) Education states</b>				
Illiterate	41		30	

Primary	26	<0.0001	34	0.1678
Secondary	18		28	
Graduate	5		18	
<b>(C) Occupation</b>				
House wife	30	<0.0001	47	<0.0001
Labourer	58		58	
Employed	2		5	
<b>(D) Residence</b>				
Rural	56	0.026	63	0.152
Urban	34		47	
<b>(E) Socio-economic class</b>				
Poor	56	<0.0001	39	0.392
Middle Class	25		41	
High Class	9		30	
<b>(F) Menstrual Period</b>				
Using Cloth	46	<0.0001	71	<0.0001
Pad	25		17	
Both	19		22	

**Table 4: Antibiotic Sensitivity Pattern of Gram Negative Bacilli (% of Sensitive isolates)**

	Imipenem	Pipracillin-Tazobactam	Amikacin	Gentamicin	Ciprofloxacin	Ofloxacin	Cefixime	Ceftriaxone
Esch. Coli	86.7	83.4	73.4	66.7	66.7	63.3	50	53
Klebsiella	87.5	87.5	75	75	62.5	62.5	58	50
Enterobacter	83.3	83.3	83.3	66.7	66.7	50	50	50
Citrobacter	100	100	75	75	50	50	50	50

## References

- UNFPA, Common Reproductive Tract Infections, No. 9,1999.
- Khan I, Khan UA. A hospital based study of frequency of aerobic pathogens in vaginal infections. J Rawal Med Coll. 2004;29(1):22–25.
- Mylonas I, Friese K. Genital discharge in women. MMW Fortschr Med. 2007;149(35-36):42–6.
- Kent HL. Epidemiology of vaginitis. Am J Obstet Gynecol 1991;165:1168–76.
- Donder GG, Vereecken A, Bosmans E, Dekeersmaecker A, Salembier G, Spitz B. Definition of a type of abnormal vaginal flora that is distinct from bacterial vaginosis: aerobic vaginitis. BJOG 2002;109: 34–43.
- Dan M, Kaneti N, Levin D, Poch F, Samra Z. Vaginitis in a gynaecologic practice in Israel: causes and risk factors. Isr Med Asso J 2003;5:629–32.
- Syed TS, Braverman PK. Vaginitis in adolescents. Adolesc Med Clin. 2004;15(2):235–51.
- Schaaf VM, Perez-Stable EJ, Borchardt K. The limited value of symptoms and signs in the diagnosis of vaginal infections. Arch Intern Med 1990, 150:1929–33.
- Bornstein J, Lakovsky Y, Lavi I, Bar-Am A, Abramovici H. The classic approach to diagnosis of vulvovaginitis: a critical analysis. Infect Dis Obstet Gynecol 2001, 9:105–11.
- Bhatia J.C., and Cleland J. "Self-reported symptoms of gynecological morbidity and their treatment in south India", Studies in family Planning, 26(4),1995,203-216.
- J.N. Krieger; et al, Diagnosis of Trichomoniasis: Comparison of conventional Met-mount examination with cytologic studies, culture and Monoclonal antibody staining of direct specimens, Journal of American Medical Association, 259(8),1998,1223-1227.
- Angelika stary, Angelika Kuchinka-koch and Lilianna Teodorowicz, Detection of Trichomonas vaginalis on Modified Columbia Agar in the Routine Laboratory, Journal of Clinical Microbiology, 40(9),2002,3277-3280.
- Thomason J.L et al, Statistical evaluation of diagnostic criteria for bacterial vaginosis, Am J Obstet Gynecol, 162(1),1990,155-160.
- King K. Holmes, Sexually transmitted diseases: Overview and Clinical approach, Harrison's Principles of internal Medicine, (New York: McGraw-Hill Professional 6<sup>th</sup> edition):768.
- Bauer AW, Kirby WM, Sherris JC and Turck M. Antibiotic susceptibility testing by standardized single disk method. Am J Clin Pathol. 1966;45:493–96.
- Clinical and Laboratory Standards Institute (CLSI) 2011; Performance Standards for Antimicrobial Susceptibility Testing; Twenty first Informational Supplement. M100-S21;31(1).
- Shamim Mumtaz, Mumtaz Ahmad\*, Irum Aftab, Naeem Akhtar, Masood ul Hassan, Abdul Hamid "Aerobic vaginal pathogens and their sensitivity pattern" J Ayub Med Coll Abbottabad 2008;20(1).
- K. M. Bansal, K. Singh, and S. Bhatnagar, "Prevalence of lowerRTI among married females in the reproductive age group (15–45)," Health and Population: Perspectives and Issues, vol. 24, no.3, pp. 157–163,2001.
- Reid G, Bruce AW. Urogenital infections in women: can probiotics help? Postgraduate Medical J 2003;79:428–32.
- Tariq N, Jaffery T, Ayub R, Alam AY, Javid MH, Shafique S. Frequency and antimicrobial susceptibility of aerobic vaginal isolates. J Coll Physicians Surg Pak 2006;16(3):196–99.

21. Mumtaz S, Akhtar N, Hayat A. Antibigram of aerobic pyogenic isolates from wounds and abscesses of patients at Rawalpindi. *Pakistan J Med Res.* 2002;41(1):16–18.
22. Vigneswaran R, McDonald PJ. Changes in vaginal flora during pregnancy and association with preterm birth. *J Inf Dis* 1994;170:724–28.
23. Akhtar N, Khan AA, Khan HH, Khan IA. Antibiotic susceptibility pattern of 196 strains of *Staph aureus* isolated from wounds and abscesses of patients at Bahawalpur. *J Ayub Med Coll Abbottabad.* 1997;9(2):29–33.
24. Hafiz S, Hafiz AN, Ali L, Chughtai AS, Memon B, Ahmad A, Hussain S, et al. Methicillin resistant *Staphylococcus aureus*; a multicentre study. *J Pak Med Assoc* 2002;52:312–5.

**How to cite this article:** Soni P, Kulsrestha S, Parihar R.S., Khatri PK. Comparative evaluation of sexually transmitted and non-sexually transmitted, reproductive tract infection at tertiary care centre in western Rajasthan, India. *Indian J Microbiol Res* 2016;3(2):107-112.