

Chronically discharging lacrimal fistula—A forerunner of Malignancy?

Babitha. V^{1,*}, Prasanna Kumari C², Raju K.V³

¹Assistant Professor, ²Associate Professor, ³Professor & HOD, Dept. of Ophthalmology, Govt. Medical College Kozhikode Kerala

***Corresponding Author:**

Email: babithavkalyan@gmail.com

Abstract

Lacrimal sac malignancies are rare cause of epiphora. Squamous cell carcinoma (SCC) is the most common type of lacrimal sac malignancy. Both sexes are equally affected^[1]. The peak incidence is at the fifth decade^[1]. About 300 cases were reported in literature^[2]. Since it is a fatal condition early diagnosis and treatment are important. We present a case of fifty nine years old chronic smoker with history of lacrimal trauma presented as lacrimal sac SCC.

Key word: Discharging fistula, Lacrimal sac malignancy, Squamous cell carcinoma

Case Report

A fifty nine year old diabetic, hyperlipidemic, chronic smoker with history of coronary artery disease presented with swelling near left medial canthus for last six years. The swelling gradually enlarged for last six months and became painful for last three months. He gave history of trauma to the left medial canthus at the age of ten with a pencil, following that he had persistent watery discharge from that area. Examination showed facial asymmetry due to periorbital oedema on left side and minimal proptosis. There was resistance to retropulsion, and positive Nafziger's sign. He had dystopia on left side, all extraocular movements were absent except for minimal abduction. FDT and FGT were positive. There was lid oedema, mechanical ptosis, conjunctival congestion on left side (Fig. 1). Bilateral cornea were clear, pupils reacting to light briskly, and BCVA of 6/9, N₆ both eyes.

Fundus was within normal limits. Supra clavicular lymphnodes were enlarged on left side. CT scan orbit showed left lacrimal sac malignancy with infiltration and extension to surrounding structures (Fig. 2). FNAC of lymphnodes showed metastasis from a poorly differentiated carcinoma possibly squamous cell carcinoma and biopsy of the lesion showed poorly differentiated squamous cell carcinoma (Fig. 3). Staging was T₄N₁M₀. Patient improved with six cycles of chemotherapy with Paclitaxel and Carboplatin. Then he lost follow up. After few months he reported with lung metastastasis. Even though two more cycles of palliative chemotherapy were given he expired with systemic metastasis.



Fig. 1: Gross appearance




Fig. 2: Histopathology of lacrimal sac malignancy



Fig. 3: Axial and coronal CT




Fig. 4

Discussion

Primary lacrimal sac tumours are rare but most of them are malignant epithelial tumours^[3,4,5]. Lacrimal sac malignancies can develop in post trauma dacryocystitis^[6]. According to Stefanyszyn MA, et al the most common clinical presentations of lacrimal drainage malignancies are epiphora, recurrent dacryocystitis, lacrimal sac mass, mass emanating through the lacrimal puncta^[2,3,4,7,8,9]. Malignant melanoma of lacrimal sac presented with bloody nasal discharge and bleeding from the punctum^[3,5,10]. The lacrimal sac malignancies were classified into epithelial and nonepithelial. Squamous and transitional cell papillomas, oncocytomas, and benign mixed tumours are benign epithelial tumours. Squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma, mucoepidermoid, adenoid cystic, and poorly differentiated carcinoma are malignant epithelial tumours. Fibrous histiocytoma, lymphoid lesions, malignant melanoma, hemangiopericytoma, lipoma, granulocytic sarcoma, neurofibroma, and very rare types like small cell carcinoma, solitary fibrous tumours, multiple myeloma, etc. are included under nonepithelial tumours^[7,11,12]. Melanoma of conjunctiva can involve lacrimal drainage apparatus^[13]. Stefanyszyn MA, et al reported that the lacrimal sac tumours had

55% malignancy rate^[7]. Squamous cell carcinoma is the most common type of epithelial malignancy^[2,4].

Dacryocystography help to diagnose the lacrimal sac tumours^[1,14]. Plain X ray may be normal or shows bone marrow erosion of lacrimal fossa and calcification in lacrimal sac malignancy^[1]. In lacrimal sac malignancies CT scan showed soft tissue mass with widening and erosion of nasolacrimal canal and inferior turbinate^[1,2,9,10,14]. MRI helps to diagnose subtle bone marrow invasion, perineural and intracranial extension^[1,2,10]. Diagnosis of lacrimal sac malignancies are by histopathological examination^[2]. Even though lacrimal drainage system malignancies are uncommon cause for lacrimal duct obstruction all DCT specimen should be sent for histopathological examination since these neoplasms are potentially life-threatening and are often difficult to diagnose^[4,10,15,16]. Malignant epithelial tumours usually recur locally and can metastasize and be fatal^[3,7,10]. The differential diagnosis of the conditions of medial canthal swellings are dacryocystitis, pseudotumour, anterior ethmoid mucocele, sarcoidosis of the sac or extensions of sinonasal malignancies^[1]. Necrotizing sialometaplasia is an inflammatory reaction against an ischemic insult or local trauma within a glandular tissue, most commonly seen in the minor salivary glands of the oral mucosa^[17]. It can rarely affect lacrimal sac mucosa (necrotizing dacryocystometaplasia)^[17]. It is considered clinically and histopathologically as a differential diagnosis of well-differentiated squamous cell carcinoma^[17].

Lacrimal sac malignancies can have lymphnode and systemic metastasis^[10]. Lacrimal sac SCC spread by direct invasion to orbit, paranasal sinuses, and skull^[1]. Lymphnode metastasis is late, involve preauricular, submandibular, jugulodigastric and cervical nodes^[1]. Effective method of managing lacrimal sac malignancy is a combined sinus orbit approach with simultaneous reconstruction of the bony defect with titanium mesh^[18]. Epithelial malignancies of the lacrimal drainage system grow along the epithelium, therefore should be treated by wide surgical excision of the tumor and the entire lacrimal drainage system combined with a lateral rhinostomy and radiation therapy^[7]. Since some of non-epithelial lacrimal sac malignancies are life threatening, early diagnosis and treatment are important^[3]. If tumor is confined to orbit at the time of diagnosis complete excision with pre and postoperative radiotherapy improves the survival^[1,10]. For tumors with orbital extension exenteration can be tried^[9]. Lacrimal sac malignancy has a recurrence rate of 50%^[2,14].

Bilateral simultaneous lacrimal sac malignancies are associated with HIV, HSV infection in young individuals^[2]. Bilateral SCC of lacrimal sac can develop denovo or seeding of tumour from other side, or as metastasis or as local recurrence, due to loss of antiangiogenic factors following excision of primary

tumour^[2]. Lacrimal sac fistula may be either congenital or acquired. Pseudo and true lacrimal fistula can develop following trauma to lacrimal region or following dacryocystitis or rupture of lacrimal abscess and can be treated by conventional DCR or endoscopic DCR^[20,21,22,23]. Literature search showed increased risk of squamous cell carcinoma in recurrent, chronic anal fistula^[24].

Conclusion

Even though benign lesions are common causes for epiphora, malignancies should be suspected especially in blood stained epiphora. Since lacrimal sac malignancies are life threatening ophthalmologist should be aware of this rare condition since early diagnosis and treatment are lifesaving. So all DCT specimen should be sent for histopathology examination. Since it has high recurrence rate follow up is mandatory.

Reference

- MR in squamous cell carcinoma of the lacrimal sac. Rahangdale SR, Castillo M, Shockley W. *AJNR Am J Neuroradiol*. 1995 Jun-Jul;16(6):1262-4.
- Bilateral squamous cell carcinoma of the lacrimal sac. *Ophthal Plast Reconstr Surg*. 2013 Nov-Dec;29(6):e149-51.. Hodgson N¹, Whipple K, Lin JH, Brumund KT, Kikkawa DO, Korn BS.
- Nonepithelial tumors of the lacrimal sac. Pe'er JJ, Stefanyszyn M, Hidayat AA. *Am J Ophthalmology*. 1994 Nov 15;118(5):650-8.
- Clinical and pathological analysis of primary lacrimal sac tumors. Bi YW, Chen RJ, Li XP. *Zhonghua Yan Ke Za Zhi*. 2007 Jun;43(6):499-504. Chinese.
- Two cases of primary malignant melanoma of the lacrimal sac. Lee HM, Kang HJ, Choi G, Chae SW, Kim CH, Hwang SJ, Lee SH. *Head Neck*. 2001 Sep;23(9):809-13.
- Epidermoid carcinoma of the lacrimalsac a fteracute posttraumatic dacryocystitis. George JL, Heymann V, Vignaud JM, Barthelme A. *J Fr Ophtalmol*. 1987;10(8-9):515-7.
- Lacrimal sac tumors. Stefanyszyn MA, Hidayat AA, Pe'er JJ, Flanagan JC. *Ophthal Plast Reconstr Surg*. 1994 Sep;10(3):169-84.
- Primary malignant melanoma of the lacrimal sac: a case report. Nam JH, Kim SM, Choi JH, Lee YK, Baek JH, Jang TJ, Park KU. *Korean J Intern Med*. 2006 Dec;21(4):248-51.
- Malignant melanoma of the lacrimal sac. Heindl LM, Schick B, Kämpgen E, Kruse FE, Holbach LM. *Ophthalmology*. 2008 Dec;105(12):1146-9. doi: 10.1007/s00347-008-1740-0.
- Primary malignant melanoma of the lacrimal sac. Li YJ, Zhu SJ, Yan H, Han J, Wang D, Xu S. *BMJ Case Rep*. 2012 Aug 13;2012.
- Neurofibroma of the lacrimal sac. Bajaj MS, Nainiwal SK, Pushker N, Sen S. *Orbit*. 2002 Sep;21(3):205-8.
- Solitary fibrous tumor of the lacrimal sac. Woo KI, Suh YL, Kim YD. *Ophthal Plast Reconstr Surg*. 1999 Nov;15(6):450-3.
- Malignant melanoma of the lacrimal drainage apparatus complicating conjunctival melanoma. Satchi K, McKelvie P, McNab AA. *Ophthal Plast Reconstr Surg*. 2015 May-Jun;31(3):207-10.
- Squamous cell carcinoma of the lacrimal drainage system: case report and literature review. De Stefani A, Lerda W, Usai A, Cavalot A, Riella P, Bussi M. *Tumori*. 1998 Jul-Aug;84(4):506-10.
- Benign mixed tumor of the lacrimal sac. Lee JS, Lee H, Chang M, Park M, Baek S. *Indian J Ophthalmology*. 2015 Mar;63(3):282-4.
- Incidence of neoplasia in patients with unilateral epiphora. Bewes T, Sacks R, Sacks PL, Chin D, Mrad N, Wilcsek G, Tumuluri K, Harvey R. *J Laryngol Otol*. 2015 Jul;129 Suppl 3:S53-7.
- Necrotizing sialometaplasia of the lacrimal sac mimicking squamous cell carcinoma: Necrotizing dacryocystometaplasia. Gomez A, Gladstone G. *Orbit*. 2016 Feb;35(1):48-50J *Laryngol Otol*. 2015 Jul;129 Suppl3:S53-7.
- En bloc resection of lacrimal sac tumors and simultaneous orbital reconstruction: surgical and functional outcomes. Alabiad CR, Weed DT, Walker TJ, Vivero R, Hobeika GA, Hatoum GF, Shriver EM, Tse DT. *Ophthal Plast Reconstr Surg*. 2014 Nov-Dec;30(6):459-67.
- 17 19Surgical treatment of mucoepidermoid carcinoma of the lacrimal sac. Yuksel D, Kosker M, Saribas F, Simsek S. *Semin Ophthalmology*. 2014 Mar;29(2):70-2.
- Pseudo external lacrimal fistula. Ittyerah TP, Fernandez ST, Joseph KA. *Indian J Ophthalmology*. 1978 Oct;26(3):21-2.
- Role of endoscopic dacryocystorhinostomy in treating acquired lacrimal fistulae. Subbaiah S. *J Laryngol Otol*. 2003 Oct;117(10):793-5.
- Acquired lacrimal sac fistula mimicking basal cell carcinoma. Litwin AS, Timlin H, Sagili S, Wright M, Malhotra R. *Br J Dermatol*. 2013 Jun;168(6):1348-50.
- Acquired lacrimal sac fistula after incision and drainage for dacryocystitis: a multicenter study. Barrett RV, Meyer DR; ASOPRS Acquired Lacrimal Fistula Study Group. *Ophthal Plast Reconstr Surg*. 2009 Nov-Dec;25(6):455-7.
- Squamous cell carcinoma arising from recurrent anal fistula. Seya T, Tanaka N, Shinji S, Yokoi K, Oguro T, Oaki Y, Ishiwata T, Naito Z, Tajiri T. *J Nippon Med Sch*. 2007 Aug;74(4):319-24.