

ANALYTICAL METHOD DEVELOPMENT AND VALIDATION FOR THE ESTIMATION OF DILTIAZEM HYDROCHLORIDE IN BULK AND PHARMACEUTICAL DOSAGE FORM BY RP- HPLC

*Amol. Y. Ghodke ¹, Bhagwat. N. Poul², Patil Bhagyashree R³

¹ Dept. of Pharmaceutical Analysis, ²Dept. of Pharmacy, ³Dept. of Quality Assurance, Maharashtra College of Pharmacy, Dept. of Pharmaceutical Analysis, Nilanga-413521 Dist. Latur Maharashtra, INDIA

*Corresponding Author:

E-mail: aamolias@gmail.com

Abstract: A simple, specific, precise and accurate Reverse Phase High Performance Liquid Chromatography (RP-HPLC) method was developed and validated for the estimation of Diltiazem HCl (Diltiazem Hydrochloride) in Bulk and Pharmaceutical dosage form. The chromatographic determination was performed on isocratic High performance liquid chromatography system of Agilent model no.1220. The separation was conducted by using column of Zorbax [C₈ (5 μ , 4.6mm \times 250)] with mobile phase consisting of buffer and Acetonitrile in the ratio of (60:40). The mobile phase was delivered at the flow rate of 1.0ml/min. The eluent was monitored at wavelength 240 nm and found a sharp and symmetrical peak with retention time of 4.66min. The method was validated for linearity, accuracy, precision, specificity, robustness. Recovery of Diltiazem HCl was found to be in the range of 98%-102%. The method was found to be linear over the concentration range 50-150 μ g/ml with coefficient correlation $r^2 = 0.995$. After developing method and studying validation parameter to obtain result complying with USP monograph.

Keywords: Diltiazem HCl, precision, accuracy, Method Validation.

Introduction

Diltiazem HCl is a calcium channel blocker, which has been used in the treatment of various cardiovascular disorders, particularly angina pectoris and systemic hypertension. Chemically it is 1, 5-Benzothiazepin-4(5H)-one, 3-(acetyloxy)-5-[2- (dimethylamino) ethyl]-2, 3-dihydro-2-(4-methoxyphenyl)-, monohydrochloride.




Fig 1: Structure of Diltiazem HCl

Literature survey revealed that methods have been developed and reported for estimation of Diltiazem HCl in bulk and in tablet dosage form and in combination also but no method was developed for estimation of Diltiazem HCl in capsule dosage form. The reported methods are HPLC and UV (2, 3, 4, 5, and 6), FT Raman spectroscopy (7), Diltiazem HCl drug with combination of other therapeutic agents HPLC and UV (8, 9, 10, and 11) and Electrophoresis method (12)

The present study was aimed to develop simple, precise, specific, accurate, linear, robust and less time consuming method.

Methods and Material

Diltiazem HCl pure drug generously gifted by local industry (Wockhardt, Aurangabad). The formulation of Diltiazem HCl capsule USP containing 90mg Diltiazem HCl was procured from local market.

Chemicals and Reagents:

Potassium monobasic phosphate (Analytical grade), Methanol (HPLC grade), Acetonitrile (HPLC grade), Triethylamine (Analytical grade), and Water (HPLC grade) were purchased from Research –Lab Fine Chem Industries, Mumbai.

Experimental Conditions:

Quantitative HPLC was performed on isocratic HPLC of Agilent model no. 1220 with ezchrom elite software G 4286B-1220 infinity isocratic LC manual injector with variable wavelength detector. For method development several trials are carried out. After many trials, the chromatographic conditions are finalised. The separation was conducted by using column of Zorbax [C₈ (5 μ , 4.6mm \times 250)] with mobile phase consisting of buffer and Acetonitrile in the ratio of (60:40). The mobile phase was delivered at the flow rate of 1.0ml/min. The eluent was monitored at wavelength 240 nm and found a sharp and symmetrical peak with retention time of 4.66min. The run time was 10 min.

Preparation of standard solution:

Transferred an accurately weighed quantity of about 60mg of Diltiazem hydrochloride into a clean dry 50ml volumetric flask, add methanol to dissolve the content and make up the volume with methanol, Dilute 1ml of resulting solution to 50ml with methanol.(12 μ g/ml)

Preparation of sample solution:

Accurately Weighed 5 Capsules and transferred the powder of capsule 213.28mg. Grind the contents thoroughly, and transfer an accurately weighed portion, equivalent to about 60 mg of Diltiazem hydrochloride, to a 100 ml volumetric flask. Add methanol and make up the volume of volumetric flask, and shake by mechanical means for 30 minutes. Sonicate the resulting solution for 10 minutes to complete the extraction. Diluting 1.12ml of resulting solution to 100 ml methanol. (12 μ g/ml)

Diluent:

The methanol is used as diluent.

Method Validation

Validation is an act of proving that any procedure, process, equipment, material, activity or system performs as expected under given set of conditions and also give the required accuracy, precision, sensitivity, etc

System suitability test:

Standard solution of Diltiazem HCl injected five times into the HPLC system. The system suitability parameters were evaluated and found to be within the limits.

Precision:

The Precision of the method was demonstrated by system precision, method precision studies and intermediate precision. In system precision, the standard solution injected five times as per procedure. In method precision six replicate injections of the standard solution and sample solution prepared as per the proposed method and chromatograms were recorded. In Intermediate precision precision was performed on different day by using different make column of same dimensions.

Accuracy:




Accuracy is calculated the percentage recovery by the assay of the known amount of analyte in the sample or as the difference between the mean and the accepted true value together with confidence intervals. For assay method, spiked samples are prepared in triplicate at three intervals over a range of 50-150% of the target concentration.

Linearity:

It is the ability of an assay to obtain test results, which are directly proportional to the concentration of an analyte in the sample. For the establishment of linearity, a minimum of 5 concentrations required. For assay of a drug substance or a finished product 50-150% of the concentration should be taken.

Specificity:

Specificity study was carried out by injecting blank, standard and sample solution of Diltiazem HCl and it shows no interference of standard and sample in the blank preparation.

**Fig 2: Chromatogram for diluent****Fig 3: Chromatogram for blank****Fig 4: Specificity Standard chromatogram of Diltiazem HCl**




Fig 5: Specificity Sample chromatogram of Diltiazem HCl

Robustness:

The robustness of the methods was determined by performing the assay of the triplicate by deliberately alternating parameters and that the results are not influenced by different changes in the above parameters.

Change in wavelength: ± 2 nm

Change in flow rate: ± 0.2 ml/min.

Change in pH: ± 0.2

Stability of analytical solution:

Evaluate the stability in analytical solution by injecting the standard preparation and sample preparation of Diltiazem HCl at regular interval. The stability of solution is carried out by 0, 3, 6, 12, 24, hrs.

Results and Discussion

The aim is to develop accurate and precise method for the quantitative estimation of Diltiazem HCl in bulk and pharmaceutical dosage form. For method development several trials are carried out. After many trials, the chromatographic conditions are finalized. The separation was conducted by using column of Zorbax [C₈ (5 μ , 4.6mm \times 250)] with mobile phase consisting of buffer and Acetonitrile in the ratio of (60:40). The mobile phase was delivered at the flow rate of 1.0ml/min. The eluent was monitored at wavelength 240 nm and found a sharp and symmetrical peak with retention time of 4.66min. The run time was 10 min.

Table 01:

Result of Assay

Sample	Amount taken	Area	
Standard	60 mg	836126	98.44%
Sample	213mg	864991	

Table 02:

Results of System suitability Test

Sr. No	Sample	Area	R.T.	Theoretical plates	Tailing factor
1.	Blank	-	-	-	-
2.	Std. 1	857171	4.643	12643	1.08
3.	Std. 2	874529	4.653	15479	0.73
4.	Std. 3	860542	4.650	12840	1.08
5.	Std. 4	884768	4.653	8631	0.82
6	Std. 5	897971	4.660	8701	0.84
Average		874996.2	4.651	11658.8	0.91
%RSD		1.93			

The blank, standard solution and sample solution are injected. There should be no interferences of standard and sample in blank preparation

Table 03:
Results of System Precision

Sr. No	Sample	Area	R.T	Theoretical plates	Tailing factor
1.	Std. 1	934671	4.627	7740	0.80
2.	Std. 2	950554	4.627	8328	0.86
3.	Std. 3	934881	4.630	8432	0.82
4.	Std. 4	956217	4.627	7653	0.81
5.	Std. 5	908143	4.627	7823	0.82
6	Std. 6	932430	4.630	7574	0.82
Average		936149	4.628	7925	0.82
% RSD		1.79			

The precision is done to check for the consistent results and which are in the limits. The method and intermediate precisions are showing the results within the limits

Table 04:
Results of Method Precision and Intermediate Precision

Sr.No	Standard (% assay)	Sample (% assay)
1	99.23	101.69
2	99.65	98.66
3	98.56	99.72
4	98.89	99.05
5	99.12	98.78
6	99.63	100.66
Average	99.18	99.76
%RSD	0.42	1.26

Table 05:
Results of specificity

Sr. No	Sample	Area	R.T	Theoretical plates	Tailing factor
1.	Blank				
2.	Std. 1	897971	4.660	8701	0.84
3.	Std. 2	900018	4.673	9011	0.87
4.	Std. 3	890884	4.680	8074	0.83
5.	Std. 4	896700	4.687	8005	0.82
6	Std. 5	894887	4.627	8421	0.82
Average		896092	4.665	8442	0.836
%RSD		0.38			

Table 06:
Results of Linearity

Concentration ($\mu\text{g/ml}$)	Peak Area
6	3119245
9	5438299
12	6925899
15	8684575
18	9735335

The proposed method is linear and the range is $6\mu\text{g/mL}$ to $18\mu\text{g/mL}$ and correlation coefficient is 0.998.




Fig 6: Graph of linearity

Table 07:
Results of Accuracy

Spike Level in %	Area	Amt. Added	Amt. Found	%recovery	Mean	SD	%RSD
50%	4142188	0.006	0.0060	100.66	98.89	1.55	1.57
	3938906	0.006	0.00585	97.70			
	4048460	0.006	0.00597	98.33			
100%	5110526	0.0012	0.00122	101.66	101.38	0.47	0.472
	5285208	0.0012	0.00121	100.83s			
	5314775	0.0012	0.00122	101.66			
150%	8651678	0.0018	0.00177	98.33	98.51	0.32	0.32
	8722465	0.0018	0.00178	98.89			
	8519110	0.0018	0.00177	98.33			

The accuracy of the method was determined by the recovery studies, carried out at different levels 50%, 100% and 150%.

Results of stability solution

Evaluate the stability in analytical solution by injecting the standard preparation and sample preparation at regular interval.

Table 08:
Results of stability solution

Stability in hours	% Assay
0	98.50
3	99.90
6	99.93
12	101.25
24	101.75
Average	100.26
%RSD	1.27

Conclusion

The developed method was found to be simple, accurate, precise, specific and robust and this method can be applied for routine quantitative analysis of Diltiazem HCl in bulk and pharmaceutical formulations like capsule.

Acknowledgements

We are grateful to Wockhardt Aurangabad India, for providing working standard. I am thankful to the my guide Dr.O.G.Bhusnure giving excellent guidance and Dr. B. N. Poul Principal of Maharashtra College of Pharmacy, Nilanga and A.Y.Ghodke Assistant professor of Maharashtra college of pharmacy Nilanga for providing necessary guidance, cooperation, facilities and chemicals, and for providing guidance on the experimental work.

References

1. United States pharmacopoeia, XXIV, United States Pharmacopoeial convention, Inc. Rockville, M.D., 2000; 573-576
2. Vivekanand A.Chatpalliwar, Pawan.K.Porwal, Neeraj Upmanyu. Validated gradient stability indicating HPLC method for determining Diltiazem hydrochloride and related substances in bulk drug and novel tablet formulation. *Journal of Pharmaceutical Analysis*; 2(3): 2012, 226–237
3. R. Pravin Cumar, Dr. M. Vasudevan, Dr. Deeca Raman. RP-HPLC Method development and validation for the estimation of Diltiazem in bulk and tablet dosage forms; *Asian Journal of Pharmaceutical and Clinical Research* , Vol 5, Issue 3, 2012, 62-64
4. H. Rowe, S. Wahab, B. Voigt. Method Development and Validation of a Stability-Indicating HPLC Assay Method for the Quantitation of Diltiazem Hydrochloride in Compounded Oral Preparations, *United States Pharmacopeia*
5. Shabana Naz, Najma Sultana, M. Saeed Arayne,Nighat Shafi. Simultaneous determination of Prazosin and Calcium chanel blockers in raw materials, pharmaceutical formulations and human serum by RP-HPLC; *International Journal of Pharmaceutical Research and Development*, VOV-2/ISSUE-9, 2010, 6-12.
6. Vidya V. Gawande and Dr. A. V. Chandewar. Development and validation of spectroscopic method for estimation of diltiazem hydrochloride in solid dosage form; *Journal of Pharmacy Research* 3(12), 2010 3032-3033
7. Jean Paul Remon, Geert J. Vergote, Chris Vervae, Tony Haemers, Francis Verpoort. Near-infrared FT-Raman spectroscopy as a rapid analytical tool for the determination of Diltiazem hydrochloride in tablets; *European Journal of Pharmaceutical Sciences* 16 (2002) 63–67
8. M.S.Bhatia, S.Kulkarni, S. D. Jadhav, S.S.Khetmar Development of Chromatographic Technique for Simultaneous Estimation of Lovastatin and Diltiazem Hydrochloride; *Mahidol University Journal of Pharmaceutical Sciences*; 39 (3-4), 2012; 17-23
9. Farhan Ahmed Siddiqui, Najma Sultana, M. Saeed Arayne, Nighat Shafi, and Azhar Hussain. Development of a RP-HPLC method for the simultaneous analysis of Diltiazem and statin: Application in pharmaceuticals and human serum; *Analytical Methods*, 2010, 2, 1571–1576 | 1571
10. Farhan Ahmed Siddiqui Najma Sultana, M. Saeed Arayne, Nighat Shafi, and Azhar Hussain. Development and Validation of New Assay Method for the Simultaneous Analysis of Diltiazem, Metformin, Pioglitazone and Rosiglitazone by RP-HPLC and applications in pharmaceuticals and human Serum; *Journal of Chromatographic Science*, Vol. 49, 2011, 774-779.
11. Chaudhari .B. G, Patel J. Development and validation of dual wavelength spectrophotometric method for simultaneous estimation of Rosuvastatin and Diltiazem in combined dosage form; *International Journal for Pharmaceutical Research Scholars*. V-1, I-3, 2012, 146-153
12. Carsten coors, Hans-George schulz, Frank Stache. Development and validation of a bioanalytical method for the quantification of Diltiazem and Desacetyl Diltiazem in plasma by capillary zone electrophoresis, *Journal of Chromatography A*, 717 (1995) 235-243.