

## Histopathological spectrum of testicular lesions- A retrospective study

Mansi Sharma<sup>1,\*</sup>, Vidhu Mahajan<sup>2</sup>, Jyotsna Suri<sup>3</sup>, KK Kaul<sup>4</sup>

<sup>1,2</sup>Senior Resident, <sup>3</sup>Professor, <sup>4</sup>Professor & HOD, Dept. of Pathology, Govt. Medical College, Jammu

**\*Corresponding Author:**

Email: drmansisharma@gmail.com

### Abstract

**Introduction & Objectives:** Testis is affected by both neoplastic and non- neoplastic conditions. Although testicular neoplasms are rare, most of the previous studies of testis are on them only. Our study was undertaken to study histopathological spectrum, age-wise distribution, laterality and clinical presentation of all testicular lesions (both neoplastic and non- neoplastic).

**Materials and Method:** This was a three year retrospective study including all the testicular specimens referred to Department of Pathology, GMC Jammu from 1st Jan 2014 to 31st Dec 2016.

**Results:** The study comprised a total of 57 cases. Non- neoplastic testicular lesions were more common than the neoplastic ones (93 vs. 7%). Non- neoplastic lesions were most common in 2nd decade of life with a wide age range of 5 months- 80 years. Among non-neoplastic lesions (n=53), undescended testis (39.62%) was the most common non-neoplastic lesion followed by inflammatory lesions (24.53%), infarcted testis (torsion, 18.86%) and atrophic testis (16.98%). Inflammatory lesions included nonspecific epididymo-orchitis (15.1%), testicular abscess (5.66%) and tubercular epididymo-orchitis (3.77%). Only 4 cases (7%) of testicular neoplasm were diagnosed in the study period amounting to only 1.33 case/ year. All 4 cases were germ cell neoplasms with age range of 14 months- 35 years and mean age 20.54 years. One case each of seminoma, yolk sac tumour, immature teratoma and mixed germ cell tumour (mixed teratoma and seminoma) was diagnosed. All the lesions were unilateral and 58% of non-neoplastic lesions were right sided. Among both neoplastic and non-neoplastic lesions, the most common symptom was testicular (scrotal/inguinoscrotal) swelling (87%). The second most common presenting complaint was empty scrotum (36.84%) and pain (36%). Inflammatory lesions in addition had history of fever(22.80%). No tumour was found in undescended testes unlike western countries.

**Conclusion:** Majority of testicular lesions are non- neoplastic and neoplastic lesions are rare, most being germ cell neoplasms. Non- neoplastic lesions are seen in all age groups but neoplasms are usually seen in younger age. Non- neoplastic lesions mimic neoplastic ones clinically, testicular swelling being the most common complaint. So histopathological examination is necessary to serve an accurate diagnosis of testicular swellings. Our findings are comparable with most studies.

**Keywords:** Undescended testis, Germ cell neoplasm, Epididymo-orchitis, seminoma, teratoma, yolk sac tumour, Non-neoplastic.

**Manuscript Received:** 30<sup>th</sup> March, 2017

**Manuscript Accept:** 29<sup>th</sup> May, 2017

### Introduction

Testis is affected by both non neoplastic and neoplastic conditions. There are various testicular lesions, ranging from paediatric to adult age groups. They usually present with scrotal swelling, pain and mass per abdomen. Non neoplastic testicular lesions include cryptorchid (undescended) testis, testicular torsion, testicular atrophy, epidermoid cysts, infections of testis like tuberculosis, infertility, malakoplakia and vasculitis.<sup>(1)</sup>

Undescended testis is the commonest genital malformation of the boys<sup>(2)</sup> and is found in approx. 1% of one year old boys.<sup>(3)</sup> An undescended testis is more likely to develop a germ cell tumour than a normally placed testis. Atrophy of testis may result from cryptorchidism, the orchitis of mumps, liver cirrhosis, estrogen administration, radiation exposure, chemotherapy, AIDS and exposure to environmental toxins.<sup>(4)</sup> Torsion of testis is a surgical emergency, commonly seen in 10-25 years of age.<sup>(5)</sup> Nonspecific epididymo-orchitis is commonly related to infections in the urinary tract and its cause varies with age. It may progress to frank abscess formation.<sup>(6)</sup> Tubercular epididymo-orchitis is a common form of genitourinary tuberculosis. It may coexist with pulmonary tuberculosis or tuberculosis of other parts of lower genitourinary system. It almost invariably begins in the epididymis and then spreads to testis. An isolated case

is rare but when it occurs, it may mimic testicular tumour.<sup>(7)</sup>

The testicular tumours although relatively rare, are of great interest and importance because of their varied histological appearances and the diverse or even conflicting views held regarding their histogenesis.<sup>(8)</sup> They account for less than 1% of all malignancies in male; constitute the 4<sup>th</sup> most common cause of death from neoplasia in a younger male, usually found in age group 15-35 years. The incidence of testicular neoplasm in western countries is rising in the past 50 years.<sup>(9)</sup> Though the etiology of testicular cancer is not well understood, various factors such as cryptorchidism, trauma, infections and genetic and endocrine factors appear to have a role in their development.<sup>(10)</sup> A definite geographic and racial distribution is seen in testicular tumours.<sup>(11)</sup> Testicular carcinoma follows a reverse pattern to most cancers with decreasing incidence rate with increasing age.<sup>(12)</sup>

Clinically the diagnosis of testicular tumours is delayed in many cases.<sup>(12)</sup> Despite new techniques in imaging and tumour marker assays, the diagnosis of testicular lesions is primarily dependent upon histopathological examination.<sup>(5)</sup> The urologists, the radiologists and chemotherapists are eventually dependent upon histological diagnosis of tumour and tumour like lesions<sup>(13)</sup> as histopathological features have

a major stake in determining the prognosis and therapeutic option.<sup>(14)</sup>

### **Aims and Objectives**

Although testicular neoplasms are rare, most of the previous studies of testis are on them only or on a single non neoplastic testicular lesion like torsion testis, undescended testis etc. Our study was undertaken to study:

- Histopathological spectrum
- Age wise distribution
- Laterality
- Clinical presentation of all testicular lesions including both neoplastic and non-neoplastic lesions in one go.

### **Materials and Method**


This was a three year retrospective study including all the testicular specimens referred to Department of Pathology, GMC Jammu from 1st Jan 2014 to 31st Dec 2016. The bilateral orchidectomy specimens of prostatic carcinoma patients, were excluded from our study. All the slides along with histopathology requisition forms containing clinical details were retrieved and reviewed. The tumours were classified according to WHO classification (2004).<sup>(15)</sup> The data collected was tabulated, analysed and compared to other similar studies.

### **Results**

- The study comprised a total of 57 cases.
- Non- neoplastic testicular lesions were more common than the neoplastic ones (93 vs 7 %, n=53 vs 4).

#### **Histopathological spectrum**


- Undescended testis (39.62%, n=21) was the most common non-neoplastic lesion followed by inflammatory lesions (24.53%, n=13), infarcted testis (torsion, 18.86%, n=10) and atrophic testis (16.98%, n=9). Inflammatory lesions included nonspecific epididymo-orchitis (15.09%, n=8), testicular abscess (5.66%, n=3) and tubercular epididymo-orchitis (3.77%, n=2) (Fig. 1).




**Fig. 1: Tubercular epididymo- orchitis (H&E, 200X)**

- Only 4 cases of testicular neoplasm were diagnosed in the study period amounting to only 1.33 case/year. All 4 cases were germ cell neoplasms. One case each of seminoma (Fig. 2), yolk sac tumour (Fig. 3), immature teratoma (Fig. 4 & 5) and mixed germ cell tumour (mixed teratoma and seminoma)


was found. No sex cord stromal tumour, lymphoma or metastasis was encountered.




**Fig. 2: Seminoma. Note lymphocytes in septae in between lobules of round uniform cells with clear cytoplasm and central nuclei. (H&E, 200X)**



**Fig. 3: Yolk sac tumour showing microcystic pattern (H&E, 200X)**



**Fig. 4: Immature teratoma showing immature neuroectodermal tissue (H&E, 200X)**



**Fig. 5: Immature teratoma showing immature cartilage (H&E, 200X)**

**Age Distribution****Table 1: Histopathological spectrum of non neoplastic testicular lesions along with age distribution**

Lesion	0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	Total
Undescended testis	4	10	4	1	1	0	1	0	21 (39.62%)
Torsion testis	0	6	1	2	1	0	0	0	10 (18.86%)
Atrophic testis	0	0	3	0	2	2	1	1	9 (16.98%)
Nonspecific epididymo-orchitis	0	0	0	1	1	4	1	1	8 (15.09%)
Testicular abscess	0	0	1	0	1	1	0	0	3 (5.66%)
Tubercular epididymo-orchitis	0	0	0	1	0	0	1	0	2 (3.77%)
Total	4	16	9	5	6	7	4	2	53 (100%)
Percentage	7.5%	30.19%	16.98%	9.43%	11.32%	13.21%	7.55%	3.77%	100%

- Non- neoplastic lesions were most common in 2nd decade of life with a wide age range of 5 months- 80 years with mean age 35.64 years. 54.71% cases were with age  $\leq 30$ .
- Undescended testis, the most common non neoplastic lesion in our study, was also most common in second decade with mean age 20.5 years and age range 5 months- 65 years. 86% were  $\leq 30$  years.
- Infarcted testis (torsion) cases ranged from 14-48 years with 70%  $\leq 30$  years and 24.1 years mean age. They were also most common in second decade.
- Atrophic testis ranged from 22-75 years with mean age 47.11 years. They were most common in third decade of life.
- Inflammatory lesions, the second most common non neoplastic lesions, ranged from 28-80 years with maximum cases in 6<sup>th</sup> decade. Mean age was 50.86 years. Among inflammatory lesions, mean ages of nonspecific epididymo-orchitis, abscess and tubercular epididymo-orchitis were 65.75, 42.33 and 51.5 years respectively.
- All 4 neoplastic cases were germ cell neoplasms with age range of 14 months- 35 years and mean age 20.54 years.

**Table 2: Age distribution of neoplastic testicular lesions**

Lesion	0-10	11-20	21-30	31-40	Total
Seminoma			1 (28 years)		1 (25%)
Immature teratoma		1 (18 years)			1 (25%)
Yolk sac tumour	1 (14 months)				1 (25%)
Mixed germ cell tumour (Mixed seminoma and teratoma)				1 (35 years)	1 (25%)
Total	1	1	1	1	4 (100%)

**Laterality**

- All the lesions were unilateral.
- 58% of non-neoplastic lesions were right sided.
- 50% of neoplastic lesions were right sided.

**Clinical Presentation**

- Among both neoplastic and non-neoplastic lesions, the most common symptom was testicular (scrotal/inguinoscrotal) swelling.
- 83.02% (n=44) of non neoplastic lesions presented with swelling with 50.94% having painless

swelling. The second most common presenting complaint was empty scrotum (39.62%) followed by pain (32.07%).

- Inflammatory lesions in addition had history of fever.
- No tumour was found in undescended testes unlike western countries.
- All 4 neoplasms presented with painless swelling.

**Discussion**

In our study the non neoplastic testicular lesions were more common than the neoplastic ones (93 vs 7%). This is in concordance with Reddy H et al (86 vs 14%)<sup>(1)</sup> and Patel MB et al (85 vs 15%)<sup>(5)</sup> but doesn't correlate with Robertson GS et al (31.5 vs 68.4%).<sup>(16)</sup>

Testicular swelling was the commonest symptom in the present study similar to previous studies<sup>(1,5)</sup> Right sided testis was more commonly involved in ours' and Patel MB's<sup>(5)</sup> studies but reverse result was found by Reddy H et al.<sup>(1)</sup>

**Table 3: Comparison of Histological Types of Non Neoplastic Testicular Lesions**

Lesion	Patel MB <sup>(5)</sup> 2015, n=85	Reddy H <sup>(1)</sup> 2016, n=86	Abba K <sup>(17)</sup> 2016, n=70	Present Study n=53
Undescended testis	8.24%	14%	10%	39.62%
Torsion testis	55.29%	22.1%	14.3%	18.86%
Atrophic testis	-	19.8%	-	16.98%
Nonspecific epididymo-orchitis	9.4%	3.5%	10%	15.1%
Testicular abscess	16.47%	19.76%	4.3%	5.66%
Tubercular epididymo-orchitis	9.4%	3.5%	12.9%	3.77%
Others	1.2%	17.44%	48.6%	-

The most common non neoplastic lesion vary from study to study. We found undescended testis and inflammatory lesions to be most common. In previous studies torsion is the most common lesion. We found non-neoplastic lesions to be most common in 2<sup>nd</sup> decade of life similar to Patel MB.<sup>(5)</sup> More than 50% were in first three decades of life in our study unlike previous studies. This variation is because of variation in relative proportion of different histological types. The age distribution of individual non neoplastic lesion was found to be in concordance with previous studies.

As described in literature, testicular tumours were found to be rare in present study also. Infact, we found them to be rarer than previous studies. We found only 4 cases amounting to only 1.33 case per year in our three year study. Incidence of testicular tumours varies from country to country and place to place thus pointing to various causative factors.<sup>(24)</sup>

**Table 4: Comparison of Average number of testicular neoplasms per year**

S No.	Study (Year)	Country	Average no. of cases per year
1	Horwich et al (2013) <sup>(18)</sup>	England	64.9
2	Walschaerts et al (2008) <sup>(19)</sup>	France	53.2
3	Chalaya PL et al (2014) <sup>(20)</sup>	Tanzania	5.6
4	Deora A et al (1994, 18 years study) <sup>(21)</sup>	India	5.5
5	Chakrabarti PR et al (2016, 10 years study) <sup>(22)</sup>	India	3.7
6	Salako AA et al (2010) <sup>(23)</sup>	South Western Nigeria	1.5
7	Present study (3 years study)	Northern India	1.33

As per Mostofi and Price,<sup>(25)</sup> germ cell tumours constitute more than 94% and stromal tumours consist of 3% of testicular tumours. Other studies also had similar results. Our 100% cases of testicular neoplasms were germ cell tumours only. No sex cord stromal tumour, lymphoma or metastasis was encountered.

**Table 5: Comparison of histological types of testicular neoplasms in various studies**

Tumour type	Patel MB <sup>(5)</sup> 2015 n=15	Deore KS <sup>(12)</sup> 2015 n=15	Chakrabarty PR <sup>(22)</sup> 2016 n=37	Reddy H <sup>(1)</sup> 2016 n=14	Sanjay M <sup>(13)</sup> 2016 n=18	Gupta A <sup>(26)</sup> 2016 n=50	Present study n=4
Seminoma	40%	26.67%	35.14%	42.9%	38.9%	48%	25%
Teratoma	33.3%	13.33%	2.7%	-	11.11%	12%	25%
Embryonal carcinoma	-	-	-	7.2%	-	16%	-
Yolk sac tumour	6.6%	6.67%	2.7%	-	5.55%	4%	25%
Mixed germ cell tumour	-	33.33%	32.43%	43%	33.33%	16%	25%
Others	20%	20%	27%	7.2%	11.11	4%	-

According to literature, the histological pattern and behaviour of the testicular tumours differs with age. Testicular neoplasm of germ cell origin is the most common malignancy in men aged between 18-35 years.<sup>(27)</sup> All our cases were also found in men younger than 40. Non seminomatous tumours are known to present in younger age than seminomatous type.<sup>(28)</sup> We had similar findings as found one pure seminoma in 3<sup>rd</sup> decade, one mixed germ cell tumour with seminoma as one component in 4<sup>th</sup> decade and two non seminomatous tumours; one yolk sac tumour and one immature teratoma in 1<sup>st</sup> and 2<sup>nd</sup> decades respectively.

Cryptorchidism is the single most important risk factor associated with testicular cancer with 10% of all testicular cancer patients having history of cryptorchidism.<sup>(20)</sup> Although we found 21 cases of undescended testis, none of them showed neoplastic focus and also none of the 4 cases of testicular neoplasms had history of undescended testis. Our finding is in concordance with Reddy H et al.<sup>(1)</sup>

Most of our findings are comparable to previous studies. Variations found may be because of small number of cases especially of tumours. A follow up study involving a larger study population and longer duration is recommended.

### Conclusion

- Majority of testicular lesions are non- neoplastic and neoplastic lesions are rare, most being germ cell neoplasms.
- Non- neoplastic lesions are seen in all age groups but neoplasms are usually seen in younger age.
- Non- neoplastic lesions mimic neoplastic ones clinically, testicular swelling being the most common complaint.
- So histopathological examination is necessary to serve an accurate diagnosis of testicular swellings.
- Our findings are comparable with most studies.

### References

1. Reddy H, Chawda H, Dombale VD. Histomorphological analysis of testicular lesions. *Indian Journal of Pathology and Oncology*, Oct-Dec 2016;3(4):558-563.
2. Mathers MJ, Sperling H, Rubben H, Roth S. The undescended testis: Diagnosis, treatment and long term consequences (Review Article). *Dtsch Arztebl Int*. 2009; 106(33):527-532.
3. Rozanski TA, Bloom D. The undescended testis: theory and management. *Urol Clin North Am* 22: 107, 1995.
4. Rosai J. Male reproductive system. In: *Rosai and Ackerman's Surgical Pathology*. 10th ed. Vol 1. Elsevier 2011: 1335-1336.
5. Patel MB, Goswami HM, Parikh UR, Mehta N. Histopathological study of testicular lesions. *Gujarat Medical Journal* 2015; 70(1):41-46.
6. Kumar V, Abbas AK, Fausto N(eds), Epstein JI. The lower urinary tract and male genital system. In: *Robbins and Cotran Pathological basis of diseases*. 7th ed. Saunders 2004:1039-41047.
7. Badmos KB. Tuberculous epididymo-orchitis mimicking a testicular tumour: a case report. *Afr Health Sci* 2012; 12(3): 395-397
8. Pratap VK, Agarwal S. Testicular neoplasms. *India Journal of cancer* 1971:40-53.
9. Bergstorm et al. Testicular cancer in nine European countries. *Int J Cancer* 1996;59:33-38.
10. Garner MJ, Turner MC, Ghadirian P, Krewski D. Epidemiology of testicular cancer: an overview. *Int J Cancer* 2005;116: 331-9.
11. Liu S, Wen SW, Mao Y, Mery L, Pouleau J. Birth cohort effects underlying the increasing testicular cancer incidence in Canada. *Can J Public Health* 1999;90:176-80.
12. Deore KS, Patel MB, Gohil RP, Delvadiya KN, Goswami HM. Histopathological analysis of testicular tumours - a 4 year experience. *International Journal of Medical Science and Public Health* 2015; Vol 4(4):554-557.
13. Sanjay M, Sushma HM. Histomorphological spectrum of tumour and tumour like lesions of testis and paratesticular structures- A cross sectional study. *Indian Journal of Pathology and Oncology*, Oct-Dec 2016;3(4):528-534.
14. Kinkade S. Testicular cancer. *Americal family Physician* 1999;59(9):2539-2550.
15. Eble JN, Sauter G, Epstein JI, et al. *Pathology and genetics of tumours of the urinary system and male genital organs*. Lyon: IARC Press; 2004.
16. Robertson GS. *Br J Surg*. 1995. Mar. 82(3): 342-5
17. Abba K, Tahir MB, Dogo HM, Nggada HA. Testicular and Paratesticular Non- Neoplastic lesions in University of Maiduguri Teaching Hospital: A 10-year Retrospective Review. *Bo Med J* 2016; 13(1): 39-44.
18. Horwich A, Nicol D, Huddart R. Testicular germ cell tumours. *BMJ*. 2013;347: 5526.
19. Walschaerts M, Huyghe E, Muller A, et al. Doubling of testicular cancer incidence rate over the last 20 years in southern France. *Cancer Causes Controls*. 2008;19: 155-61.
20. Chalya PL, Simbila S, Rambau PF. Ten- year experience with testicular cancer at a tertiary care hospital in a resource-limited setting: a single centre experience in Tanzania. *World Journal of Surgical Oncology*. 2014; 12: 356-63.
21. Deotra A, Mathur DR Vyas MC. A 18 years study of testicular tumour in Jodhpur, western Rajasthan. *Ind. Journal of surgery* 1994;40(2):68-70.
22. Chakrabarti PR, Dosi S, Varma A, Kiyawat P, Khare G, Matreja S. Histopathological trends of testicular neoplasm: An experience over a decade in a tertiary care centre in the Malwa belt of central India. *Journal of Clinical and Diagnostic Research*. 2016 Jun, Vol-10(6): EC16-EC18.
23. Salako AA, Onakpoya UU, Osasan SA, Omoniyi-Esan GO. Testicular and paratesticular tumours in south western Nigeria. *Afr Health Sci*. 2010; 10: 14-17.
24. K.P. Dieckmann and U. Pichlmeier. Clinical epidemiology of testicular germ cell tumours. *World Journal of Urology*. April 2004 Vol.22, Issue I: 2-14.
25. Mostofi FK, Price EB, Jr. Tumors of the male genital system. *Atlas of Tumor Pathology, Fascicle 7, Series 2*. Washington, DC: Armed Forces Institute of Pathology, 1973. Pp. 1186-1200.
26. Gupta A, Gupta S, Gupta S, Gupta V. Testicular tumours: A histopathological study of 50 cases. *Indian Journal of Pathology and Oncology*, Oct-Dec 2016; 3(4):544-547.
27. Shanmugalingam T, Soultati A, Chowdhury S, Rudman S, Van Hemelrijck M. Global incidence and outcome of testicular cancer. *Clin Epidemiol*. 2013; 5: 417-27.
28. Ruf CG, Isbaran H, Wagner W, et al. Changes in the epidemiological features of testicular germ cell cancer: age at diagnosis and relative frequency of seminomas are constantly and significantly increasing. *Urol Oncol*. 2014;32:33.